

Angular momentum I ground state probabilities of boson systems interacting by random interactions

Y. M. Zhao^{a,b}, A. Arima^c, and N. Yoshinaga^a,

^a Department of Physics, Saitama University, Saitama-shi, Saitama 338 Japan

^b Department of Physics, Southeast University, Nanjing 210018 China

^c The House of Councilors, 2-1-1 Nagatacho, Chiyodaku, Tokyo 100-8962, Japan

In this paper we report our systematic calculations of angular momentum I ground state probabilities ($P(I)$) of boson systems with spin l in the presence of random two-body interactions. It is found that the $P(0)$ dominance is usually not true for a system with an odd number of bosons, while it is valid for an even number of bosons, which indicates that the $P(0)$ dominance is partly connected to the even number of identical particles. It is also noticed that the $P(I_{max})$'s of bosons with spin l do not follow the $1/N$ ($N = l + 1$, referring to the number of independent two-body matrix elements) relation. The properties of the $P(I)$'s obtained in boson systems with spin l are discussed.

PACS number: 21.60.Ev, 21.60.Fw, 05.30.Jp, 24.60.Lz

It was discovered a few years ago [1] that the dominance of the 0^+ ground state (0 g.s.) of even fermion systems is obtained by diagonalizing a scalar two-body Hamiltonian which is randomly determined. Many works have been done in the context of this discovery [2-16]. For example, many efforts were made concerning the origin of this observation and generic properties of many-body systems in the presence of random interactions.

In the case of d -boson systems the 0 g.s. probability (denoted as $P(0)$) was found to be close to 0% periodically when the number of d bosons $n = 6k \pm 1$, where k is a positive integer [10]. This finding of systematic counter examples of 0 g.s. dominance in d -boson systems suggests that the breaking of the 0 g.s. dominance might be related to certain features associated with bosonic degrees of freedom and thus could be easily seen also for bosons with other l . To see whether this is true or not we report in this paper systematic calculations of boson systems with spin l in the presence of random interactions.

The Hamiltonian that we take is as follows:

$$H_d = \sum_l \frac{1}{2} \sqrt{2l+1} G_L \left((b_l^\dagger \times b_l^\dagger)^{(L)} \times (\tilde{b}_l \times \tilde{b}_l)^{(L)} \right)^{(0)} \quad (1)$$

where $b_l^\dagger$ is the creation operator of a boson with spin l , and $\tilde{b}_l$ is the time-reversal of the annihilation operator b_l . The G_L 's are two-body interactions between the bosons, and are assumed to follow a two-body random ensemble (TBRE), i.e., the G_L 's are a set of random numbers with a distribution function

$$\rho(G_L) = \frac{1}{\sqrt{2\pi}} \exp(-G_L^2/2), \quad L = 0, 2, \dots, 2l. \quad (2)$$

All the results in this paper are obtained by using 1000 sets of the above two-body random interactions.

We first look at the case of four bosons with spin l . These systems are interesting and worthwhile to study separately, because their fermionic counterparts, namely, four fermions in a single- j shell, were studied extensively [7, 10, 11] since these systems are the simplest and non-trivial ones to investigate the 0 g.s. dominance of fermions. Fig. 1 shows a few important angular momentum I ground state probabilities versus l for systems with boson number $n=4$. One sees that the pattern of I g.s. probabilities ($P(I)$'s) of four

bosons with spin l is very similar to that of four fermions in a single- j shell. For instance, $P(0)$'s vs. l stagger with an interval of $\delta_l = 3$; $P(0)$'s are dominant over other $P(I)$'s except for only two cases: $l = 2$ and 8 ; $P(I_{max})$'s decrease with l . A new feature is that the $P(l)$'s are considerably large (small) when l is even (odd), i.e., the $P(l)$'s exhibit an even-odd staggering behavior. We should be aware that $I = l$ is possible in the four boson case, while in the case of four fermions there are no $I=j$ states.

The 0 g.s. dominance in the case of four bosons with spin l can be understood based on the following observation: there is *only* one non-zero eigenvalue (denoted as $E_0^{L(l)}$) among all eigenvalues of $I = 0$ states if only one of $G_L = -1$ and others are zero. This observation was proved recently by constructing the $I = 0$ states using a pair basis [16]. This means that there is a very large probability for $I = 0$ states to give the lowest eigenvalue, compared to other I states for which there are many non-zero eigenvalues. According to our empirical rule [10], $P(I) = \mathcal{N}_I/N$, where $\mathcal{N}_I$ is the number of times that the ground state has angular momentum I when one of the two-body matrix elements is -1 and others are zero, and N is the number of two-body matrix elements. The $P(0)$'s are therefore expected to be larger than all the other $P(I)$'s (with only two exceptions of $l = 2$ and 8).

In Fig. 1 one sees that the increase of $P(0)$'s “coincides” with that of $D_0^{(l)}$, the number of $I = 0$ states of four bosons with spin l . $D_0^{(l)}$ is 1, 1, 1, 2, 2, 2, 3, 3, 3, 4, 4, 4, $\dots$ for $l=0, 1, 2, 3, \dots$, etc. Although we still can not prove this correlation, below we give an argument based on our empirical rule [10]. From the sum rule of diagonal matrix elements [17], one obtains that $\sum_L E_0^{L(l)} = -\frac{1}{2}n(n-1)D_0^{(l)} = -6D_0^{(l)}$, where n is the number of bosons in the system and $n = 4$ here. This relation means that the sum of $E_0^{L(l)} \times (-1)$ increases with $D_0^{(l)}$, suggesting that $\mathcal{N}_0$ increases with $D_0^{(l)}$ simultaneously. Because $P(0) = \mathcal{N}_0/N$, a regular increase of $D_0^{(l)}$ of four bosons with spin l [18] produces a regular staggering of the $P(0)$'s. It is noted that a similar “coincidence” of the $P(0)$'s staggering with increasing number of $I=0$ states was firstly observed in [7, 10] for four fermions in a single- j shell, but without an explanation or argument.

We next go to systems with arbitrary particle number n , in order to investigate general

features of the $P(I)$'s of boson systems with spin l in the presence of random interactions. Here a particularly interesting question should be asked concerning the $P(I)$'s for an odd number of particles: which angular momentum I g.s. dominance will appear, $I = 0$ or $I = l$? In systems with an odd number of fermions in a single- j shell, there is no $I=0$ state. One expects $I = j$ g.s. probabilities to be large, because one easily relates large j g.s. probabilities to large 0 g.s. probabilities for the neighboring even n , and the calculations using different types of random interactions showed [7, 10] that this is indeed the case. For systems with odd n , however, one may have $I=0$ states and thus it is unknown a priori whether the $I = 0$ or $I = l$ g.s. dominance occurs in these systems.

Fig. 2 shows the $P(l)$'s and $P(0)$'s in boson systems with $l=4$ and 6, and n running as large as possible. For odd n and $l = 1$ or $n = 3$ and any odd l , there is no $I = 0$ state; for odd n and $l = 3, 5, 7$ and 9, the $I = 0$ states do not exist unless $n \geq 15, 9, 7$, and 5, respectively. For fifteen bosons with $l = 3$, nine bosons with $l = 5$, seven bosons with $l = 7$ and five bosons with $l = 9$ or 11, $P(0) \sim 0\%$ always according to our calculations using 1000 sets of a TBRE Hamiltonian. From Fig. 2 and these odd- l cases we conclude that the $P(0)$'s are in general much less than the corresponding $P(l)$'s when n is odd. On the other hand, the $P(0)$'s are mostly larger than the $P(l)$'s when n is even. This result indicates that the 0 g.s. dominance is robust for systems with even n , but not true generally if n is odd. In the latter case, we observe that the 0 g.s. dominance is easily lost, with a few exceptions. We therefore expect that the 0 g.s. dominance is partly connected to the even number of particles.

It was shown in previous calculations that the 0 g.s. dominance may appear in boson systems with odd n , such as sp [8, 9], sd [2, 9, 10], and sdg [10] boson systems. However, to a large extent, the very large $P(0)$ therein are associated with the s boson condensation, which contributes around 40% of $P(0)$ [9]. In other words, sp and sd systems are very special systems in which the s boson condensation produces 0 g.s. dominance when n is odd. Without s bosons, the $P(0)$'s of those systems with odd n would be drastically smaller than those with s bosons, and g.s. probabilities for other I 's, such as $I = l$ or $I = I_{max}$, would be much larger. It would be very interesting to carry out systematic

calculations of $P(I)$'s for both even and odd numbers of bosons with many l 's.

Now we come to the I_{max} g.s. probabilities of bosons with spin l interacting by a TBRE Hamiltonian. It was shown in [10] that the $P(I_{max})$'s are associated with the number of independent two-body matrix elements, and not sensitive to the particle numbers of the system. For fermions in a single- j shell, $P(I_{max}) \sim 1/N$, where $N = j + \frac{1}{2}$. This relation is also applicable to sd and sdg boson systems. Thus it is interesting to check whether the $P(I_{max})$'s of more general boson systems vs. l show a similar pattern or not.

Fig. 3 shows the $P(I_{max})$'s vs. l with n ranging from three to six. One sees large deviations of the $P(I_{max})$'s calculated by diagonalizing a TBRE Hamiltonian from those predicted by the $1/N = 1/(l + 1)$ relation. The $P(I_{max})$'s are systematically larger than $1/N$ (when spin l is small the agreement is rather good, however), and increase with n .

An argument why the behavior of the $P(I_{max})$'s is different for bosons and fermions is as follows. It is known from previous studies [10] that the large $P(I_{max})$ comes from a gap produced by the pairing interaction $G_{L_{max}}$ for bosons with spin l , and the pairing interaction $G_{J_{max}}$ for fermions in a single- j shell, where $L_{max} = 2l$ and $J_{max} = 2j - 1$, respectively. For both cases one may evaluate the gap associated with $G_{J_{max}}$ or $G_{L_{max}}$ using analytical formulas of $(E_{I_{max}-2} - E_{I_{max}})$, because here the state with I_{max} ($I_{max} - 2$) is found to be the ground (first excited) state if $G_{J_{max}}$ or $G_{L_{max}}$ is -1 . We obtain $(E_{I_{max}-2} - E_{I_{max}})$ as follows:

$$\begin{aligned}
\text{boson systems : } & \frac{2ln - 1}{4l - 1}; \\
\text{fermion systems, } n = 4 : & \frac{3}{8} + \frac{105}{108(4j - 7)} + \frac{135}{64(4j - 5)} \\
& + \frac{63}{128(4j - 3)}; \\
\text{fermion systems, } n = 5 : & \frac{35}{128} + \frac{2205}{2048(4j - 9)} + \frac{5145}{2048(4j - 7)} \\
& - \frac{1785}{2048(4j - 5)} - \frac{189}{2048(4j - 3)}; \\
\text{fermion systems, } n = 6 : & \frac{27}{128} + \frac{10395}{8192(4j - 11)} + \frac{2835}{1024(4j - 9)}
\end{aligned}$$

$$-\frac{4725}{4096(4j-7)} - \frac{45}{256(4j-5)} - \frac{297}{8192(4j-3)}. \quad (3)$$

One easily sees that the gap for bosons with spin l increases regularly with n at an interval $\sim 1/2$ if l is large, while that for fermions in a single- j shell is much smaller (almost one order) in magnitude and comparable for different n and j . For instance, the gap is 0.47, 0.39, 0.35 for $n=4, 5$ and 6 fermions in a $j=15/2$ shell, respectively, while the gap is 2.03, 2.56, 3.07 for $n=4, 5, 6$ bosons with spin $l=7$, respectively. According to the empirical rule [10], a relatively larger gap makes the corresponding $P(I)$ larger: the larger the gap is, the larger the corresponding $P(I)$ is. Therefore, the $P(I_{max})$'s of bosons with spin l can be expected to be larger than $1/N = 1/(l+1)$, which works well for fermions and some boson systems with small l 's studied in [10], such as d , sd and sdg bosons.

To summarize, in this paper we reported systematic calculations of angular momentum I ground state probabilities for systems of bosons with spin l in the presence of random interactions. First, we found that the 0 g.s. probabilities $P(0)$'s of four bosons with spin l exhibit a pattern similar to that of four fermions in a single- j shell. We explained the 0 g.s. dominance and the staggering of the $P(0)$'s based on a special property of $I=0$ states when only one of the G_L 's are -1 and the other two-body matrix elements are zero.

Second, we found that the 0 g.s. dominance is usually not true for systems with an odd number of bosons with spin l (though the $I=0$ states may exist), which indicates that the 0 g.s. dominance is partly connected to the even number of particles.

Third, we found that the I_{max} g.s. probabilities of bosons with spin l are systematically larger than those predicted by the $1/N$ relation ($N=l+1$ for bosons with spin l), and that $P(I_{max})$ increases with boson number n . We presented an argument for this difference between bosons and fermions in terms of a large gap for bosons systems with G_{2l} being -1 and others zero.

One of the authors (ZHAO) would like to thank the Japan Society for the Promotion of Science (ID: P01021) for supporting the present work.

References

- [1] C. W. Johnson, G. F. Bertsch, D. J. Dean, Phys. Rev. Lett. **80**, 2749(1998); C. W. Johnson, G. F. Bertsch, D. J. Dean, and I. Talmi, Phys. Rev. **C61**, 014311(1999).
- [2] R. Bijker and A. Frank, Phys. Rev. Lett. **84**, 420(2000); Phys. Rev. **C62**, 14303(2000).
- [3] D. Kusnezov, N. V. Zamfir, and R. F. Casten, Phys. Rev. Lett. **85**, 1396(2000).
- [4] M. Horoi, B. A. Brown, and V. Zelevinsky, Phys. Rev. Lett. **87**, 062501(2001).
- [5] V. Velazquez and A. P. Zuker, Phys. Rev. Lett. **88**, 072502 (2002); Y. M. Zhao and A. Arima, nucl-th/0202056.
- [6] R. Bijker, A. Frank, and S. Pittel, Phys. Rev. **C60**, 021302(1999).
- [7] D. Mulhall, A. Volya, and V. Zelevinsky, Phys. Rev. Lett. **85**, 4016(2000); Nucl. Phys. **A682**, 229c(2001).
- [8] D. Kusnezov, Phys. Rev. Lett. **85**, 3773(2000); *ibid.* **87**, 029202 (2001); R. Bijker and A. Frank, Phys. Rev. Lett. **87**, 029201(2001).
- [9] R. Bijker and A. Frank, Phys. Rev. **C64**, 061303 (2001); *ibid.* **C65**, 044316.
- [10] Y. M. Zhao, A. Arima, and N. Yoshinaga, Phys. Rev. **C66**, 034302 (2002); *ibid.* **C66**, 064322 (2002).
- [11] Y.M. Zhao and A. Arima, Phys. Rev. **C64**, (R) 041301(2001).
- [12] Y. M. Zhao, S. Pittel, R. Bijker, A. Frank, and A. Arima, Phys. Rev. **C66**, (R) 041301 (2002); Y. M. Zhao, A. Arima, and N. Yoshinaga, *ibid.* **C66**, 064323 (2002).
- [13] Lev Kaplan, Thomas Papenbrock, and Calvin W. Johnson, Phys. Rev. **C63**, 014307(2001).
- [14] P. H-T. Chau, A. Frank, N. A. Smirnova, and P. V. Isacker, Phys. Rev. **C66**, (R) 061302 (2002).


- [15] V. Velazquez, J. G. Hirsch, A. Frank, and A. P. Zuker, Phys. Rev. **C67**, 034311(2003).
 - [16] Y. M. Zhao, A. Arima, J. N. Ginocchio, and N. Yoshinaga, (to be published).
 - [17] N. Yoshinaga, A. Arima, and Y. M. Zhao, J. Phys. **A35**, 8575(2002).
 - [18] J. N. Ginocchio and W. C. Haxton, *Symmetries in Sciences VI*, Edited by B. Gruber and M. Ramek (Plenum Press, New York, 1993), p. 263; Y. M. Zhao and A. Arima, (to be published).
-

Fig. 1 The I g.s. probabilities vs. l of four bosons with spin l . The results are obtained by 1000 runs of a TBRE Hamiltonian, except that we present the $P(0)$'s (open squares) predicted by a prescription given in Ref. [10], showing that the approach of Ref.[10] is reasonably applicable to bosons with spin l .

Fig. 2 The $I = 0$ and $I = l$ g.s. probabilities versus n of $l = 4$ and 6. One sees that $P(0)$ is in general smaller than the corresponding $P(l)$ when n is an odd number, indicating that the 0 g.s. dominance is partly associated with the even number of particles.

Fig. 3. The $P(I_{max})$'s vs. l with n ranging from three to six. It is seen that the $P(I_{max})$'s are systematically larger than $1/(l+1)$, and increase with n but seem to saturate around $n = 6$.

Fig. 1






Fig. 2 January 7th/2003

